

⑫

**EUROPEAN PATENT APPLICATION**

⑳ Application number: 88308024.4

㉑ Int. Cl.4: **A61K 31/71 , C07H 17/00**

㉒ Date of filing: 31.08.88

㉓ Priority: 10.09.87 PCT/US87/02317

㉔ Date of publication of application:  
15.03.89 Bulletin 89/11

㉕ Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

㉖ Applicant: **PFIZER INC.**  
235 East 42nd Street  
New York, N.Y. 10017(US)

㉗ Inventor: **Remington, Jack Samuel**  
1995 Val Pariaso Avenue  
Menlo Park California(US)

㉘ Representative: **Moore, James William, Dr.**  
Pfizer Limited Ramsgate Road  
Sandwich Kent CT13 9NJ(GB)

㉙ **Azithromycin and derivatives as antiprotozoal agents.**

㉚ The use of azithromycin, its 4"-epimer, or the 4"-amino-4"-deoxy analogs thereof in the treatment of systemic protozoal infections in mammals, particularly toxoplasmosis in man; and pharmaceutical compositions therefor.

**EP 0 307 128 A2**

## AZITHROMYCIN AND DERIVATIVES AS ANTIPROTOZOAL AGENTS


Background of the Invention

5 The present invention is directed to the use of compounds of the formula (I) as defined below, viz., azithromycin, its 4"-epimer, and corresponding 4"-deoxy-4"-amino analogs in the treatment of systemic protozoal infections in mammals, particularly in the treatment of toxoplasmosis, a protozoal infection due to strains of Toxoplasma gondii, particularly troublesome in pregnant women and among those such as AIDS patients, who are immune deficient.

10 Azithromycin is the U.S.A.N. (generic name) for 9a-aza-9a-methyl-9-deoxo-9a-homoerythromycin A, a broad spectrum antibacterial compound derived from erythromycin A. Azithromycin was independently discovered by Bright, U.S. Patent 4,474,768 and Kobrehel et al., U.S. Patent 4,517,359. The name "N-methyl-11-aza-10-deoxo-10-dihydroerythromycin A" was employed in these patents. The present more systematic name is based upon the ring expansion and replacement nomenclature of the "IUPAC  
15 Nomenclature of Organic Chemistry, 1979 Edition," Pergamon Press, 1979, pp. 68-70, 459, 500-503. 4"-Epi-azithromycin (4"-epi-9a-aza-9a-methyl-9-deoxo-9a-homoerythromycin A), 4"-amino-4"-deoxy-azithromycin (4"-amino-9a-aza-9a-methyl-9-deoxo-4"-deoxy-9a-homoerythromycin A), and 4"-epi-4"-amino-4"-deoxyazithromycin A (4"-epi-4"-amino-9a-aza-9a-methyl-9-deoxo-4"-deoxy-9a-homoerythromycin A), also broad spectrum antibacterials derived from erythromycin A, are the subjects of Bright, U.S. Patent  
20 4,526,889, Hauske and Nagel, U.S. Patent 4,512,982, and Hauske and Nagel, loc. cit., respectively.

There is a continuing need for drugs which are effective against protozoal infections in mammals, in particular against toxoplasmosis in man. Transmission of the disease may occur transplacentally, by ingestion of raw or undercooked meat containing tissue cysts, or by exposure to oocysts in cat feces. Neonatal congenital toxoplasmosis, which is acquired transplacentally, the mother having acquired  
25 primary infection during or prior to pregnancy, can lead to spontaneous abortion, miscarriage or still-birth, birth defects, or the birth of a child with the clinical disease. The disease can cause brain damage and even death in those having weakened immune systems, particularly among those suffering from AIDS (acquired immune deficiency syndrome) where toxoplasma encephalitis is a commonly found, life threatening infection. Heretofore, there has been no alternative to the present regimen of pyrimethamine plus a  
30 sulfonamide - a relatively toxic regimen with numerous side effects among the latter patient population. Approximately 20% of AIDS patients are seropositive for Toxoplasma antibodies and approximately 30% of these seropositive individuals will suffer toxoplasmic encephalitis, reflecting the critical problem in this patient population. In one recent series, approximately 50% of the patients died, median time to death being 4 months. Furthermore, since the incidence of relapse is also prohibitively high, new drugs are  
35 needed which can be given both for initial treatment and as suppressive therapy for the life of the patient.

It has recently been reported that the macrolide antibiotic, roxithromycin (the 9-[O-(2-methoxyethoxymethyl)]oxime of erythromycin A) possesses activity against toxoplasmosis in mice (see Hofflin and Remington, Antimicrobial Agents and Chemotherapy, vol. 31, pp. 346-348 (1987); and leading references there cited).



(la)  $R^1 = \text{OH}$ ,  $R^2 = \text{H}$  azithromycin

(lb)  $R^1 = \text{H}$ ,  $R^2 = \text{OH}$  4'-epi-azithromycin

(lc)  $R^1 = \text{NH}_2$ ,  $R^2 = \text{H}$  4'-amino-4'-deoxy-azithromycin

(ld)  $R^1 = \text{H}$ ,  $R^2 = \text{NH}_2$  4'-epi-4'-amino-4'-deoxy-azithromycin

### Summary of the Invention

We have now found that the compounds of the formula (I), wherein one of  $R^1$  and  $R^2$  is hydrogen and the other is hydroxy or amino (conveniently named herein as azithromycin derivatives, *vide supra*) possess remarkably potent activity against Protozoa, particularly Toxoplasma species, and so are valuable in pharmaceutical compositions for a method of treating or preventing protozoal infections in mammals, including man. These compounds are especially valuable in the treatment of toxoplasmosis, an infection due to a strain of Toxoplasma gondii, which, as noted above, is a particular problem in pregnant women and in immune compromised patients.

### Detailed Description of the Invention

The present invention is readily carried out. The compounds of the formula (I) are prepared according to the methods of U.S. Patents 4,474,768, 4,512,982 and 4,526,889, cited above, which are hereby included by reference. A particularly valuable form of azithromycin (la) for this purpose is azithromycin dihydrate prepared according to methods disclosed in Examples below.

The utility of the compounds of the formula (I) in the treatment or prevention of protozoal infections in mammals is demonstrated by their remarkable activity in model Toxoplasma gondii infections in mice. For example, we have found azithromycin (la) to have potent *in vivo* activity against murine toxoplasmosis. Mice infected intraperitoneally with  $10^2$  tachyzoites of the virulent RH strain of T. gondii and treated 24 hours later with 200 mg azithromycin/kg/day orally by gavage (solubilized in polyethylene glycol 200) for 10 days all survived. Concentrations of 100 or 50 mg/kg resulted in 80 and 20% survival, respectively. Further experiments revealed that one daily dose of 200 mg/kg for each of 3 days after infection resulted in 100% survival of mice infected with  $10^3$  RH tachyzoites. Moreover, this concentration of the drug protected 100% of infected mice when administered as late as 72 hours after infection with  $10^2$  RH tachyzoites. Additional experiments revealed that 70% of mice infected intracerebrally with  $10^4$  tachyzoites of the C56 strain of T. gondii and treated with 200 mg/kg/day for 10 days survived, but only 10% of untreated controls survived. These results indicate that azithromycin is highly effective against infection with T. gondii. (See Hofflin et al., cited above, and references there cited, for more detailed descriptions of these murine toxoplasmosis models).

In the treatment or prevention of systemic protozoal infections in mammals, particularly toxoplasmosis in man due to strains of Toxoplasma gondii, the compounds of the formula (I), including the pharmaceutically acceptable salts thereof, are dosed orally or parenterally. Oral dosage will generally be preferred, particularly in cases where the drug is dosed chronically as a preventive measure. However, particularly in acute administration for severe cases of toxoplasmosis, parenteral administration may be preferred, a matter to be determined at the discretion of the attending physician. The preferred dosage range is about 5-100 mg per kg of body weight per day, in single or divided daily doses, regardless of the route of administration. In special situations, particularly in life-threatening cases of infection, higher doses may be prescribed at the discretion of the attending physician.

When used to treat or prevent a systemic protozoal infection in a mammal, particularly toxoplasmosis in man, the compounds of the formula (I), including the pharmaceutically acceptable salts thereof, can be dosed alone, but are preferably dosed in the form of pharmaceutical compositions comprising the active compound and a pharmaceutically-acceptable carrier or diluent. Such pharmaceutical compositions, whether for oral or parenteral administration, are prepared according to conventional methods of pharmacy, for example, as disclosed in U.S. Patents 4,474,768, 4,512,982 and 4,526,889, cited above, and included by reference.

The present invention is illustrated by the following example, but is not limited to the details thereof.

### EXAMPLE 1

#### Non-Hygroscopic Azithromycin Dihydrate

##### Method A

The hygroscopic monohydrate of Preparation 1 (100 g; water-content:3.1%), tetrahydrofuran (220 ml) and diatomaceous earth (5 g) were combined in a 500 ml Erlenmyer flask, stirred for 30 minutes and filtered with 20 ml of tetrahydrofuran wash. The combined filtrate and wash was transferred to a 3 liter round bottom flask. The solution was stirred vigorously and H<sub>2</sub>O (2.0 ml) was added. After 5 minutes, hexane (1800 ml) was added over 5 minutes, with continued vigorous stirring. Following an 18 hour granulation period, title product was recovered by filtration with 1 x 10 ml hexane wash, and dried in vacuo to 4.6±0.2% H<sub>2</sub>O by Karl Fischer. 89.5 g.

##### Method B

The hygroscopic monohydrate of Preparation 1 (197.6 g) and tetrahydrofuran (430 ml) were charged to a reactor and the mixture stirred to achieve a milky white solution. Activated carbon (10 g) and diatomaceous earth (10 g) were added and the mixture stirred for 15 minutes, then diluted with 800 ml of hexane and filtered with suction over a pad of diatomaceous earth with 250 ml of hexane for wash. The combined filtrate and wash was diluted to 2500 ml with hexane and warmed to 34 °C. With stirring, 24.7 ml of H<sub>2</sub>O was added. The mixture was allowed to cool to room temperature, granulated for five hours and title product recovered and dried as in Method A. 177.8 g.

The dihydrate melts sharply at 126 °C (hot stage, 10 °C/minute); differential scanning calorimetry (heating rate, 20 °C/minute) shows an endotherm at 127 °C; thermal gravimetric analysis (heating rate 30 °C/minute) shows a 1.8% weight loss at 100 °C and a 4.3% weight loss at 150 °C; ir (KBr) 3953, 3553, 3488 2968, 2930, 2888, 2872, 2827, 2780, 2089, 1722, 1664, 1468, 1426, 1380, 1359, 1344, 1326, 1318, 1282, 1270, 1252, 1187, 1167, 1157, 1123, 1107, 1082, 1050, 1004, 993, 977, 955, 930, 902, 986, 879, 864, 833, 803, 794, 775, 756, 729, 694, 671, 661, 637, 598, 571, 526, 495, 459, 399, 374, 321 and 207 cm<sup>-1</sup>; [alpha]<sub>D</sub><sup>26</sup> = -41.4 ° (c = 1, CHCl<sub>3</sub>).

Anal. Calcd. for C<sub>38</sub>H<sub>72</sub>N<sub>2</sub>O<sub>1.2</sub>·2H<sub>2</sub>O:  
C, 58.14; H, 9.77; N, 3.57; OCH<sub>3</sub>, 3.95; H<sub>2</sub>O, 4.59.  
Found:

C, 58.62; H, 9.66; N, 3.56; OCH<sub>3</sub>, 4.11; H<sub>2</sub>O, 4.49.

Neutralization Equivalent (0.5N HCl in 1:1 CH<sub>3</sub>CN:H<sub>2</sub>O):

Calcd.: 374.5. Found: 393.4.

Samples of a dihydrate, slightly over dried to contain 4.1% water (less than theoretical) rapidly picked-up water at 33%, 75% or 100% relative humidities to achieve the theoretical water content (4.6%) for the dihydrate. At 33% and 75% relative humidities, water content remained essentially constant for at least 4 days. At 100% relative humidity, the water content further rose to about 5.2, where it remained essentially constant of the next three days.

A sample of the same dihydrate, maintained at 18% relative humidity gradually lost water. At four days, the water content was 2.5% and at 12 days, 1.1%.

## EXAMPLE 2

15

### Azithromycin Powder for Oral Suspension

20 The following powdered ingredients were thoroughly blended:

Azithromycin Dihydrate (1200 g on anhydrous basis) 1268.5 g

Sucrose 23000 g

Sodium phosphate tribasic dodecahydrate 250 g

Sodium benzoate 90 g

25 Hydroxypropylcellulose 40 g

Xanthan gum 40 g

Certified food coloring agent(s) in solid form 3 g or as required to achieve the desired color

Fruit and/or vanilla flavoring agents in solid form 440 g or as required to achieve the desired taste

The resulting blend contains 47.75 mg of azithromycin activity per gram. Amber screw cap bottles (60 ml) are filled with 10.47 g of the blend. Prior to oral administration as a suspension, distilled water is added (25 ml) and the mixture shaken. One teaspoon (5 cc) of this mixture provides a 100 mg dose of azithromycin. Higher or lower doses are achieved by appropriate modification of the dosage volume.

35

## EXAMPLE 3

### Azithromycin Capsules (250 mg) for Oral Administration

40

The following ingredients were accurately weighed, combined, and blended in a suitable blender for 15 minutes.

Hydrated azithromycin 3360.9 g\*

45 (3250.0 g on an anhydrous basis)

Anhydrous lactose 2015.9 g

Corn starch 611.0 g

The blended material was milled through Fitz JT mill with a No. 2A plate (0.093") at slow speed with knives forward, the milled mixture blended for an additional 15 minutes, and weighed. The resulting milled and blended mixture (5977.2 g) was then blended for 5 minutes with a 9:1 lubricant mixture of magnesium stearate:sodium lauryl sulfate (91.65 g), the further blend slugged on a Stokes DD-2 fitted with six stations of 3/4" flat faced punches, and the slugs granulated by remilling and additionally blending as specified above. Additional 9:1 lubricant (29.5 g) was blended with the resulting granulated blend (5869 g) and the material encapsulated into #0 capsules on a Zanasi RM-63 capsule machine at a fill weight of 483±23 mg to yield capsules containing no more than 275 mg and no less than the desired 250 mg of azithromycin activity.

By appropriately modifying the capsule size, the fill weight and the proportion of azithromycin in the

blend, capsules containing 100 mg, 125 mg, 375 mg or 500 mg of azithromycin activity are prepared.

4'-Epi-azithromycin, 4'-amino-4'-deoxy-azithromycin and 4'-epi-4'-amino-4'-deoxy-azithromycin capsules are prepared in like manner, substituting in equal weight of the active ingredient (corrected for potency as free base) for azithromycin.

5

#### EXAMPLE 4

10

#### Azithromycin Tablets (250 Mg) for Oral Administration

The following ingredients were accurately weighed, combined and blended in a suitable blender for 30 minutes:

15 Azithromycin dihydrate 14245.0 g\*  
(13,485.0 g on an anhydrous basis)  
Dibasic calcium phosphate 22205.0 g  
AC-DI-SOL 1620.0 g  
20 Magnesium stearate 1242.7 g

The blend was milled in a Fitzpatrick D comminutor fitted with a No. 3 plate (0.125") with knives forward at 3600 rpm, then blended for an additional 30 minutes. To the resulting milled blend (39,192 g) was added an additional 783.8 g of magnesium stearate and blending continued for 5 minutes. The mixture was then  
25 slugged according to the preceding example, and remilled as immediately above, and blended for 5 minutes. Additional magnesium stearate (394.5 g) was added to the resulting granulated blend (39,445 g), blending was continued for 5 minutes, and the mixture tableted on a Killian tableting machine with forced feeder and 32" x 5/8" upper and lower oval shaped punches, each tablet having a weight of 787 mg  $\pm$  37 mg, each containing no less than 250 mg and no more than 275 mg of azithromycin activity.

30

#### EXAMPLE 5

35

#### Azithromycin for I.V. or I.M. Injection

In a sterile environment and using sterile, particle free equipment and components, 10,949 g of water  
40 for injection was placed in a compounding flask. Anhydrous citric acid, 494.4 g was added and dissolved with agitation. In a separate flask 310 g of sodium hydroxide was dissolved in 690 g of water. A portion of the latter (755 g) was used to adjust the pH of the citric acid from 1.63 to  $5.09 \pm 0.02$ . Azithromycin dihydrate 670.0 g (equivalent to 642.5 g of anhydrous base) was added, and the mixture adjusted to pH  $6.60 \pm 0.1$  with 4.0 g additional of the sodium hydroxide solution. Water (6076.5 g) was added to bring the  
45 resulting solution to a final weight of 18,948.9 g. If desired, the solution is sterile filtered at this stage, using a millipore filter. Using a filling machine, 50 ml flint type vials were each filled with  $15.06 \pm 0.45$  g of this solution, loosely stoppered with gray teflon stoppers, and freeze dried to yield stoppered vials each containing  $51 \pm 1.5$  mg of azithromycin activity in the form of freeze dried solids. Prior to i.m. or i.v. injection, water for injection (10 ml) is added by injection by syringe through the stopper, and the freeze  
50 dried solids redissolved by shaking. Virtually the entire contents of the vial is taken up into the syringe and injected either i.v. or i.m.

55

#### PREPARATION 1

Hygroscopic Azithromycin Monohydrate

Substantially following the methylation procedure of Kobrehel et al., U.S. Patent 4,517,359; and the crystallization procedure of Bright, U.S. Patent 4,474,768: 9-deoxy-9a-aza-9a-homoerythromycin A (previously called 11-aza-10-deoxy-10-dihydroerythromycin A; 100 g, 0.218 mol) was dissolved with stirring in 400 ml  $\text{CHCl}_3$ . Formic acid (98%; 10.4 ml, 0.436 mol) and formaldehyde (37%; 16.4 ml, 0.349 mol) were added over 4-5 minutes, and the mixture heated at reflux for 20 hours. The mixture was cooled to ambient temperature, diluted with 400 ml  $\text{H}_2\text{O}$  and adjusted to pH 10.5 with 50% NaOH. The aqueous layer was separated and extracted 2 x 100 ml with fresh  $\text{CHCl}_3$ . The organic layers were combined, stripped in vacuo to 350 ml, twice diluted with 450 ml of ethanol and restripped to 350 ml, and finally diluted with 1000 ml  $\text{H}_2\text{O}$  over a 1 hour period, pausing for 15 minutes as a slurry began to develop after the addition of about 250 ml of  $\text{H}_2\text{O}$ . Title product was recovered by filtration and dried in air at 50°C for 24 hours. 85 g; mp 136°C; differential thermal analysis (heating rate 20°C/minute) shows an endotherm at 142°C; thermal gravimetric analysis (heating rate 30°C/minute) shows a 2.6% weight loss at 100°C and a 4.5% weight loss at 150°C; water content 3.92%; ethanol content 1.09%.

Anal. Calcd. for  $\text{C}_{38}\text{H}_{72}\text{N}_2\text{O}_{12}$  (corrected for ethanol and water content):

C, 58.46; H, 9.78; N, 3.74; Alkoxy, 4.67.

Found: C, 58.40; H, 9.29; N, 3.50; Alkoxy, 4.52.

A sample of the monohydrate (having a water content of 3.2%) was maintained at 18% relative humidity for 14 days. The sample lost water over the first 24 hours to yield monohydrate having the theoretical water content (2.35%). The water content then remained substantially constant over 14 days, a value of 2.26% being recorded at 14 days.

At 33% relative humidity the water content of a sample of the same monohydrate rapidly rose to 5.6% where it remained substantially steady for at least three days. Similarly at 75% and 100% relative humidity, the water content rose rapidly, but was now maintained at even higher levels, 6.6% and 7.2%, respectively, for at least 3 days.

### 30 Claims

1. Use of an antiprotozoal effective amount of:

azithromycin;

4"-epi-azithromycin;

35 4"-amino-4"-deoxy-azithromycin; or

4"-epi-4"-amino-4"-deoxy-azithromycin;

or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for use in a method for the treatment or prevention of a protozoal infection in a mammal.

2. Use of an anti-Toxoplasma gondii effective amount of:

40 azithromycin;

4"-epi-azithromycin;

4"-amino-4"-deoxy-azithromycin; or

4"-epi-4"-amino-4"-deoxy-azithromycin;

or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for use in a method for the treatment or prevention of an infection in a mammal due to a strain of Toxoplasma gondii.

3. The use of a compound according to claim 1 or claim 2 wherein said compound is azithromycin or a pharmaceutically acceptable salt thereof.

4. The use of a compound according to claim 1 or claim 2 wherein said compound is 4"-epi-azithromycin or a pharmaceutically acceptable salt thereof.

50 5. The use of a compound according to claim 1 or claim 2 wherein said compound is 4"-amino-4"-deoxy-azithromycin or a pharmaceutically acceptable salt thereof.

6. The use of a compound according to claim 1 or claim 2 wherein said compound is 4"-epi-4"-amino-4"-deoxy-azithromycin or a pharmaceutically acceptable salt thereof.

**THIS PAGE BLANK (USPTO)**



(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 307 128  
A3**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: **88308024.4**

(51) Int. Cl.<sup>5</sup>: **A61K 31/71, C07H 17/00**

(22) Date of filing: **31.08.88**

(30) Priority: **10.09.87 PCT/US87/02317**

(43) Date of publication of application:  
**15.03.89 Bulletin 89/11**

(84) Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

(88) Date of deferred publication of the search report:  
**04.07.90 Bulletin 90/27**

(71) Applicant: **PFIZER INC.**  
**235 East 42nd Street**  
**New York, N.Y. 10017(US)**

(72) Inventor: **Remington, Jack Samuel**  
**1995 Val Paraiso Avenue**  
**Menlo Park California(US)**

(74) Representative: **Moore, James William, Dr.**  
**Pfizer Limited Ramsgate Road**  
**Sandwich Kent CT13 9NJ(GB)**

(54) **Azithromycin and derivatives as antiprotozoal agents.**

(57) The use of azithromycin, its 4"-epimer, or the 4"-amino-4"-deoxy analogs thereof in the treatment of systemic protozoal infections in mammals, particularly toxoplasmosis in man; and pharmaceutical compositions therefor.

**EP 0 307 128 A3**

Xerox Copy Centre



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
P,X	ANTIMICROBIAL AGENTS AND CHEMOTHERAPY, vol. 32, no. 5, May 1988, pages 755-757, American Society for Microbiology; F.G. ARAUJO et al.: "Azithromycin, a macrolide antibiotic with potent activity against toxoplasma gondii" * Whole article * ---	1-6	A 61 K 31/71 C 07 H 17/00
P,X	ANTIMICROBIAL AGENTS AND CHEMOTHERAPY, vol. 32, no. 4, April 1988, Pages 524-529, American Society for Microbiology; H.R. CHANG et al.: "In vitro effects of four macrolides (Roxithromycin, Spiramycin, Azithromycin [CP-62,993], and A-56268) on toxoplasma gondii" * Whole article * ---	1-6	
P,X	AIDS, vol. 2, suppl 1, 1988, pages S177-S182, Gower Academic Journals Ltd; A. PITCHENIK: "The treatment and prevention of mycobacterial disease in patients with HIV infection" * Page S179 * ---	1-6	TECHNICAL FIELDS SEARCHED (Int. Cl.4)  A 61 K C 07 H
D,Y	EP-A-0 101 186 (PFIZER INC.) * Whole document * ---	1-6	
D,Y	EP-A-0 109 253 (PFIZER INC.) * Whole document * ---	1-6	
		-/-	
The present search report has been drawn up for all claims			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>12-03-1990</b>	Examiner <b>FOERSTER W.K.</b>
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document  T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ***** & : member of the same patent family, corresponding document			

EPO FORM 1500 03.82 (P0401)



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
D, Y	ANTIMICROBIAL AGENTS AND CHEMOTHERAPY, vol. 31, no. 2, February 1987, pages 345-348, American Society for Microbiology; J.M. HOFFLIN et al.: "In vivo synergism of Roxithromycin (RU 965) and interferon against Toxoplasma gondii" * Whole article *	1-6	
Y	ANTIMICROBIAL AGENTS AND CHEMOTHERAPY, vol. 31, no. 5, May 1987, pages 811-812, American Society for Microbiology; M. WALSH et al: "In viro evaluation of CP-62,993, Erythromycin, Clindamycin, and Tetracycline against Chlamydia trachomatis" * Whole document *	1-6	
A	EUR: J. CLIN. MICROBIOL., vol. 6, no. 1, February 1987, pages 1-2; F. ARAUJO et al.: "Toxoplasmosis in immunocompromised patients"		
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
Place of search THE HAGUE		Date of completion of the search 12-03-1990	Examiner FOERSTER W.K.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document	

EPO FORM 1503 (03.82) (P0401)

**THIS PAGE BLANK (USPTO)**